

Euthanasia Without Venous Access; Intraorgan injections you can do with clients present

Kathleen Cooney, DVM, CHPV, DACAW resident
Companion Animal Euthanasia Training Academy
Loveland, Colorado

Learning Objectives:

1. Effectively administer euthanasia solution using each technique
2. Make better decisions towards technique selection
3. Appreciate the complexities of the procedure

A Need for Greater Understanding

While euthanasia is considered a necessary part of veterinary medicine, the skill set needed to proficiently perform it is often overlooked by teaching institutions. Only a small number of veterinary institutions devote lecture time to describe the euthanasia procedure in detail. Most of the time, the euthanasia discussion pertains to the emotional impact of such a decision rather than the medical procedure itself. To be successful, veterinary professionals must understand all that goes into the euthanasia experience including meeting the client's needs and the pharmacology/physiology behind euthanasia.

When euthanasia is necessary, skilled personnel are required to perform the procedure to ensure death is achieved in accordance with the American Veterinary Medical Association's (AVMA) Guidelines for the Euthanasia of Animals. As of 2019, the guidelines state, "Each facility or institution where euthanasia is performed is responsible for training its personnel adequately to ensure it operates in compliance with federal, state, and local laws. Training and experience should include a familiarity with the normal behavior and anatomy of the species being euthanized and an understanding of the mechanism by which the selected technique induces loss of consciousness and death." Poorly performed euthanasias complicate the mourning process. With proper education and training in euthanasia, the incidence of bad euthanasias, or dysthanasias, will decrease which is essential for the pet and client. There is no reason for a client to experience a technically challenging euthanasia because of a lack in training by their veterinarian and staff. In 2016, the Companion Animal Euthanasia Training Academy (CAETA) was founded as a training resource for veterinarians, technicians, and anyone wanting to learn best practices.

Choosing the Right Technique

When choosing a euthanasia technique, veterinarians must choose one that is appropriate for the situation and take all factors into consideration. The method of euthanasia you choose will depend on many things:

1. Comfort with the technique

2. Supplies

3. The presence of onlookers
4. The type and amount of euthanasia solution available
5. The signalment and physical condition of the animal

Also critical to consider early on in choosing a technique are the AVMA's Criteria for Euthanasia. Every technique a veterinarian might use must be checked against these 14 criteria. Not everyone can be met, but they should at least be considered as the technique is chosen. It is important the client understands why a particular technique is chosen and exactly what to expect as death occurs. Their physical, spiritual, and emotional needs should be addressed so that the experience is as enriching as possible. Details such as where to gather, who should be there, aftercare arrangements, etc. should all be discussed before the time for euthanasia arrives. The use of pre-euthanasia sedatives and anesthesia should also be discussed. More and more families are requesting their use to help ensure a peaceful passing. Planning ahead will eliminate fear and help loved ones move forward in a healthier manner.

With any of these techniques, take your time and be consistent. If you feel the method you start out with is not going to work, readily move on to another to ensure things continue smoothly for the client and the pet. You will have to decide just how fast to move onto another technique if the first is not working.

The following list of techniques are AVMA approved and are considered common techniques for dogs, cats, and some exotics. Remember that intraorgan injections in the US require patient unconsciousness which can be facilitated with anesthetic drugs. Drug volumes are based on euthanasia solutions of 390 mg/ml concentration and lethal mammalian doses are included for each technique according to the AVMA.

Intravenous Injection (85 mg/kg)

Pros

Fast and effective
Standard dosing of euthanasia solution
Sedation not required

Cons

Venous access necessary
Blood pressure concerns
Requires moderate skill

To perform an intravenous injection, you need to find a vein to work with. The accessory cephalic and cephalic veins in the front leg and the medial and lateral saphenous veins in the back leg are most common. Each has its pros and cons. Pick the one that is most appropriate under the circumstances.

Prepare your solution ahead of time if you can. IV injections using a barbiturate require dosing of 1ml per 10 pounds of body weight. If your euthanasia solution is extremely thick, you can dilute it with saline to ease its movement through the needle and catheter. The saline does not negatively affect cardiac arrest time or postmortem side effects. If you insert your euthanasia syringe with needle directly into the vein, make sure to draw back and check for blood. To be

safe, you should draw back at least once during venous administration to make sure that your needle is still placed correctly.

Once you have your needle inserted directly into the vein or a catheter placed, inject slowly and steadily to avoid putting too much pressure on your set up. When all of the solution has been injected, and you are certain it all went into the vein, you can remove your needle. Apply pressure to your injection site to stop any bleeding. If you have a catheter placed, you can flush with saline to clear your catheter of the euthanasia solution. When performing an intravenous injection, remember that death occurs very quickly. You should start to see the effects of your injection within 30 seconds or so. Let the client know what they might see before you start injecting. Once death has occurred, remove the catheter and wipe your injection site with the hydrogen peroxide-soaked gauze. Note that catheter placement is not required for IV administration, but the AVMA advises their use when owners are present.

Intracardiac Injection (85 mg/kg)

Pros

Fast and effective
Eliminates venous pressure concerns
Works for all domesticated pets

Cons

General perception as gruesome
Requires advanced skill
Requires sedation

The easiest way to give an intracardiac injection is to have the pet lying in lateral recumbency on either their right or left side. I prefer to have a dog lying on its right side because the heart is easier to auscultate from the left and the left ventricle is usually the easiest chamber to inject. The heart in most dogs and cats will reside from the 2nd or 3rd intercostal space (ICS) to the 5th or 6th ICS and from the sternum to about two-thirds of the way up the thorax. In my experience, the heart is usually more cranial and ventral than I think it will be. When ausculting the heart, you need to pinpoint the PMI or Point of Maximum Intensity. On the left side of the chest, this will likely be the point of the aortic valve located in the 4th intercostal space at the level of the shoulder. On the right side, the right AV valve will be the loudest and is also in the 4th intercostal space at the level of the olecranon/elbow. The olecranon is located near the 5th intercostal space so there will be little heart caudal to this point. Use your stethoscope or even your hand to find the Point of Maximum intensity (PMI) on the chest wall. Then, grasp the lower antebrachium and press the elbow up the chest wall to simulate where it would normally be if the pet were standing. A good place to insert your needle is usually just a bit cranial to the point of the elbow. Combining the location of the PMI with this landmark should help you find the heart.

Your patient must be fully sedated and unconscious. It should not react to your injection in any way. Make sure you draw up enough euthanasia solution to stop the heart. You don't want to have to inject a second time unless absolutely necessary. If your patient weighs 40 pounds, draw up at least 6 mls. This way you have a little extra to spare in case you accidentally inject some outside of the heart. Use a larger syringe than is necessary to hold the solution. When injecting into the heart, you have to draw back blood to make sure you are within a chamber

and therefore you need a little extra room in your syringe. If giving 6 mls of solution, I will use a 12 ml syringe. If there is pulmonary effusion present, you might miss the heart on your first stick and accidentally draw up some of the fluid. If you have extra room in the syringe, you can gently redirect and try again without having to draw up a new syringe full of euthanasia solution. If you find yourself within an airway, you will draw back air, and again it is nice to have a little extra space available in your syringe.

When performing an intracardiac injection in larger dogs, you will need to use a long needle. My service uses 1.5" to 2" 18 gauge needles for dogs over 20 pounds. As you prepare to enter the chest wall, keep your needle perpendicular to the body. Angling the needle will increase the distance you need to travel before hitting the heart. If you contact a rib during penetration of the chest wall, either start over or gently walk your needle tip off the rib edge and keep going. When you think you are in the heart, draw back on the plunger. If you draw back and get negative pressure, your needle tip is within something solid (e.g., the myocardium, a tumor, etc.). Push the needle in farther if you can and try to draw back again. If you still do not aspirate blood, gently redirect and try again, without removing the needle completely from the chest. Remember that if you let go of a syringe that is inserted into the heart, it may move along with the contractions. Once you see blood in your syringe, you are free to administer your solution.

Intraperitoneal Injection (255 mg/kg)

Pros

Easy to perform

Works in all domesticated pets

No sedation required*

Cons

Longer time to absorb

More solution needed

Possible to accidentally inject organs

While not technically an intraorgan injection, we include it here for completion. An intraperitoneal injection means that the euthanasia solution is given into the peritoneal space, i.e. the abdomen. The solution must therefore miss neighboring organs or the administration would be considered intraorgan, and without sedation, is considered painful. Even though pre-sedation is not required for this technique, I always give it. *Sedation is required if using a class III euthanasia solution.

Two areas of injection are the ventral midline caudal to the xiphoid process and low on the right lateral abdomen. The needle should be inserted at an angle slightly toward the head and the syringe plunger pulled to aspirate for negative pressure or air. If no blood or fluid is seen in the syringe, you are free to inject. Because the euthanasia solution is moving into the blood stream through absorption across abdominal organ membranes and serosal linings, it may take longer to achieve cardiac death. Three times the IV dose is required for this technique. If your pet is not pre-sedated and is still conscious after 20-30 minutes, you will need to give more solution. When necessary, massage the abdomen to help the solution absorb.

Intrahepatic Injection (170 mg/kg)

Pros

Simple technique
Works well when other techniques fail
Works in all companion pets

Cons

Requires deep sedation or anesthesia
More solution needed

The liver is large, highly vascular, and is usually easy to palpate. Veterinarians choose intrahepatic injections over intraabdominal injections because of the improved uptake of the euthanasia solution. When an intravenous injection is not viable, an intrahepatic injection can be a great alternative. Like intracardiac injections, intrahepatic injections need to be done under deep sedation or anesthesia.

Choose a needle length that is sufficient for reaching the liver. With small pets, a 1-inch needle should be adequate, but in larger pets, a 1.5" or 2" needle may be necessary. I also recommend giving more euthanasia solution than you would use for an intravenous euthanasia. I generally double the dose because the rate of absorption can be delayed based on how well you place your injection. If you miss the liver, the death may not occur in a reasonable amount of time so having more solution present will help speed things along.

To inject the liver, place your needle in the notch on either side of the xyphoid process and aim cranially about 45 degrees, up under the last rib of your laterally recumbent pet. You can even apply some inward pressure with the syringe or with your free hand to allow the needle to move deeper. There is no need to draw back and check for blood. Keep the needle in the ventral half of the abdomen to avoid the stomach, which is located more dorsally. I like intrahepatic injections because they work well from either the left or right side. Death should occur within about two minutes or so with a well-performed intrahepatic injection, but I usually tell families that their pet may pass immediately or it may take a few minutes.

Intrarenal Injections (255 mg/kg)

Pros

Great for cats and small mammals
Fast and effective

Cons

Requires deep sedation or anesthesia
Requires moderate skill

This method is a great choice if venous access is difficult or when preparing a catheter site is too obtrusive. As with intrahepatic injections, the kidneys will speed the rate of absorption over standard intraabdominal injections. Like with cardiac and hepatic injections, deep sedation or anesthesia is required. I will use cats as my example in describing how to perform an intrarenal injection.

When choosing which kidney to inject, pick the one that you can feel and isolate the best. For me, this is almost always the left one. You should be able to grasp it easily and curl your fingers around it. If a cat is in renal failure, the kidneys may be smaller, have a nodular feeling, or even be impossible to locate. If you have doubts about one kidney, consider using the other. Also

make sure that you are not palpating a fecal ball in the colon. If you are using the right kidney, you may have to use your forefinger to manipulate it into reach. Gently push your finger up under the last rib to find it and move it caudally. It will only shift by a centimeter or two.

Once your feline patient is deeply asleep, you can let the owner hold the cat on his or her lap or you can place the cat on any flat surface. Gently run your hands along the abdomen to find the kidney and feel for any abdominal muscle tensing. If the cat tenses, it is not sedated enough for you to proceed. If necessary, give more sedation until no response is noted. When you are ready to inject, use the hand you will use to inject to push the kidney upward, from the cat's downside, into your holding hand and cup the kidney under your fingertips raising it up parallel with the spine. If you have a hard time doing this, just hold it firmly in a position and don't let it slip out of your grasp. You want to keep hold of the kidney through your entire injection. If you accidentally let go mid-injection, try to hold it in place within the abdomen using your fingers and finish the injection. If you cannot find the kidney again, gently advance the syringe toward the liver and finish your injection.

As is the case with intrahepatic injections, you need to give more solution than an intravenous injection would require. Standard injection amount is three times the IV dose. My service has a standard protocol of 6 mls per renal injection in a cat of any size. This helps ensure euthanasia takes place quickly and standardizes our record keeping.

Upon injection, you should feel the kidney swell in your fingers. This is a good indication that you are positioned properly. Continue to inject all of your solution. If you are certain you are positioned correctly and the kidney does not swell, you might be injecting into the renal pelvis, which may slow absorption. Kidney swelling does not guarantee immediate death, but it does increase the odds that death will occur faster. Either way, watch the cat for cessation of breathing. Most of the time, your feline patient will stop breathing even before you finish the injection. As with intrahepatic injections, let the client know ahead of time that their cat may pass immediately or within just a few minutes.

Switching from Plan A to Plan B

When preparing to euthanize a patient, one must not only determine the best technique to use but have a backup technique in mind should things get tricky. When your patient presents to you, it's wise to consider first which sedatives or anesthetics will induce the best sleep, and based on this, determine which euthanasia technique will lead to the shortest Time To Death (TTD). Keep in mind that fast death does not equal the best death, but for the most part, fast is what clients expect. Let's take the example of a 30kg bulldog with peripheral edema. If Plan A is to give euthanasia solution via the vein, it is best to administer pre-euthanasia drugs that will maintain strong blood pressures for venous access. If a vein cannot be located, Plan B for a dog of this size would be an intracardiac injection, followed by Plan C's intrahepatic injection. Before even beginning the procedure, the proper supplies must be acquired including enough solution to efficiently bring about death. Switching to another method should be done quickly and without too much discussion with the clients. As long as the veterinary team is conveying

confidence and control over the situation, they will trust them to make the right decision for their pet.

Summary

Euthanasia, quality euthanasia, is an art form all its own. We need to fully understand the nuances of each available method to know how best to utilize it when the time comes. It's common to have a preferred technique based on the species and even breed of pet. Pre-euthanasia sedation and anesthesia is now commonplace in many countries, especially for those hospitals and mobile services specializing in enhanced end-of-life experiences. Intraorgan injections are rapidly becoming as common as intravenous injections. Veterinary teams are encouraged to learn more about the techniques discussed here and bolster confidence in their ability to reach for another, less familiar technique should the need arise. Please consider learning more with the Companion Animal Euthanasia Training Academy (CAETA) and other dedicated euthanasia organizations.

References

Leary S, Pharmaceuticals F, Underwood W, et al. AVMA Guidelines for the Euthanasia of Animals: 2020 Edition. 2020.

Morris P. Blue Juice: Euthanasia in Veterinary Medicine. Temple University Press; 2012. Available at: <https://www.jstor.org/stable/j.ctt14bt7d8>. Accessed April 8, 2023.

Robertson SA. Pharmacologic Methods: An Update on Optimal Pre-sedation and Euthanasia Solution Administration. *Vet Clin North Am Small Anim Pract* 2020;50:525–543.

Cooney K. Common and Alternative Routes of Euthanasia Solution Administration. *Vet Clin North Am Small Anim Pract* 2020;50.

Orvin L. Case Reports: Challenging Euthanasia Cases. *Vet Clin North Am Small Anim Pract* 2020;50:647–652.

Cooney, KA. et al. *Veterinary euthanasia techniques : a practical guide*. 2012. Ames, Iowa: Wiley-Blackwell.

Cooney, KA et al. The use of intrarenal injection of sodium pentobarbital for euthanasia in cats. *American journal of veterinary research*. 2022. [Online] 83 (1), 95–99.